

## Bibliographie

- Basey, A. C., Fant, J. B. et Kramer, A. T. (2015). Producing native plant materials for restoration: 10 rules to collect and maintain genetic diversity. *Native Plants Journal*, 16(1), 37-53. <https://doi.org/10.3368/npj.16.1.37>
- Bower, A. D., Clair, J. B. St. et Erickson, V. (2014). Generalized provisional seed zones for native plants. *Ecological Applications*, 24(5), 913-919. <https://doi.org/10.1890/13-0285.1>
- Breton, G., Guêné-Nanchen, M. et Rochefort, L. (2026). Long-term assessment of the Moss Layer Transfer Technique for the restoration of Sphagnum-dominated peatlands. *Restoration Ecology*, 34(1), e70212. <https://doi.org/10.1111/rec.70212>
- Brown, S. C. et Bedford, B. L. (1997). Restoration of wetland vegetation with transplanted wetland soil: An experimental study. *Wetlands*, 17(3), 424-437. <https://doi.org/10.1007/BF03161432>
- Budelsky, R. A. et Galatowitsch, S. M. (2000). Effects of water regime and competition on the establishment of a native sedge in restored wetlands. *Journal of Applied Ecology*, 37(6), 971-985. <https://doi.org/10.1046/j.1365-2664.2000.00540.x>
- Buisson, E., Jaunatre, R., Römermann, C., Bulot, A. et Dutoit, T. (2018). Species transfer via topsoil translocation: lessons from two large Mediterranean restoration projects. *Restoration Ecology*, 26(S2). <https://doi.org/10.1111/rec.12682>

- Bullock, J. M. (1998). Community translocation in Britain: Setting objectives and measuring consequences. *Biological Conservation*, 84(3), 199-214.  
[https://doi.org/10.1016/S0006-3207\(97\)00140-7](https://doi.org/10.1016/S0006-3207(97)00140-7)
- Chenot, J., Jaunatre, R., Buisson, E. et Dutoit, T. (2017). Long-term effects of topsoil transfer assessed thirty years after rehabilitation of dry alluvial quarries in Southeastern France. *Ecological Engineering*, 99, 1-12.  
<https://doi.org/10.1016/j.ecoleng.2016.11.010>
- Craft, C. (2022). *Creating and restoring wetlands: From theory to practice (2nd edition)*. Elsevier. <https://doi.org/10.1016/B978-0-12-823981-0.00013-7>
- Ding, X., Su, P., Zhou, Z. et Shi, R. (2019). Belowground Bud Bank Distribution and Aboveground Community Characteristics along Different Moisture Gradients of Alpine Meadow in the Zoige Plateau, China. *Sustainability*, 11(9), 2602.  
<https://doi.org/10.3390/su11092602>
- Flora of North America Editorial Committee. (2000). *Flora of North America*, Vol. 22 Magnoliophyta: Alismatidae, Arecidae, Commelinidae (in part), and Zingiberidae. Oxford University Press. Oxford, United Kingdom, 352 p.
- Fowler, W. M., Fontaine, J. B., Enright, N. J. et Veber, W. P. (2015). Evaluating restoration potential of transferred topsoil. *Applied Vegetation Science*, 18(3), 379-390.  
<https://doi.org/10.1111/avsc.12162>
- Gerrits, G. M., Waenink, R., Aradottir, A. L., Buisson, E., Dutoit, T., Ferreira, M. C., Fontaine, J. B., Jaunatre, R., Kardol, P., Loeb, R., Magro Ruiz, S., Maltz, M., Pärtel, M., Peco, B., Piqueray, J., Pilon, N. A. L., Santa-Regina, I., Schmidt, K. T., Sengl, P.,

- ... Wubs, E. R. J. (2023). Synthesis on the effectiveness of soil translocation for plant community restoration. *Journal of Applied Ecology*, 60(4), 714-724.  
<https://doi.org/10.1111/1365-2664.14364>
- Holl, K. D., Luong, J. C. et Brancalion, P. H. S. (2022). Overcoming biotic homogenization in ecological restoration. *Trends in Ecology & Evolution*, 37(9), 777-788.  
<https://doi.org/10.1016/j.tree.2022.05.002>
- Hugron, S., Guêné-Nanchen, M., Roux, N., LeBlanc, M.-C. et Rochefort, L. (2020). Plant reintroduction in restored peatlands: 80% successfully transferred – Does the remaining 20% matter? *Global Ecology and Conservation*, 22, e01000.  
<https://doi.org/10.1016/j.gecco.2020.e01000>
- Kiehl, K., Kirmer, A., Donath, T. W., Rasran, L. et Hölzel, N. (2010). Species introduction in restoration projects – Evaluation of different techniques for the establishment of semi-natural grasslands in Central and Northwestern Europe. *Basic and Applied Ecology*, 11(4), 285-299. <https://doi.org/10.1016/j.baae.2009.12.004>
- Larson, M. A. & Titus, J. E. (2018). Urban wetland seed bank profiles in south-central New York State<sup>1, 2</sup>. *The Journal of the Torrey Botanical Society*, 145(2), 126.  
<https://doi.org/10.3159/TORREY-D-17-00034.1>
- Liu, Q., Jiang, M., Lyu, X. & Wang, G. (2014). A Review of Similarity between Soil Seed Bank and Aboveground Vegetation in Wetlands. *Acta Ecologica Sinica*, 34(24).  
DOI:10.5846/stxb201303120403

Mahaney, W. M., Wardrop, D. H. et Brooks, R. P. (2005). Impacts of sedimentation and nitrogen enrichment on wetland plant community development. *Plant Ecology*, 175(2), 227-243. <https://doi.org/10.1007/s11258-005-0011-2>

McKay, J. K., Christian, C. E., Harrison, S. & Rice, K. J. (2005). "How Local Is Local?" – A Review of Practical and Conceptual Issues in the Genetics of Restoration. *Restoration Ecology*, 13(3), 432–440. <https://doi.org/10.1111/j.1526-100X.2005.00058.x>

McKnight, S. K. (1992). Transplanted seed bank response to drawdown time in a created wetland in East Texas. *Wetlands*, 12(2), 79-90. <https://doi.org/10.1007/BF03160589>

Natural Resources Conservation Service. (2025). *PLANTS Database*. United States Department of Agriculture. Accessed October 13, 2025. <https://plants.usda.gov>

Phartyal, S. S., Rosbakh, S., Ritz, C. & Poschlod, P. (2020). Ready for change: Seed traits contribute to the high adaptability of mudflat species to their unpredictable habitat. *Journal of Vegetation Science*, 31(2), 331–342. <https://doi.org/10.1111/jvs.12841>

Purre, A.-H., Truus, L. et Ilomets, M. (2021). A Decade of Vegetation Development on Two Revegetated Milled Peatlands With Different Trophic Status. *Mires and Peat*, 27. <https://doi.org/10.19189/MaP.2019.BG.StA.1928>

- Sparks, E. L., Cebrian, J., Biber, P. D., Sheehan, K. L. et Tobias, C. R. (2013). Cost-effectiveness of two small-scale salt marsh restoration designs. *Ecological Engineering*, 53, 250-256. <https://doi.org/10.1016/j.ecoleng.2012.12.053>
- Streever, B. et Zedler, J. (2000). To plant or not to plant. *BioScience*, 50(3), 188. [https://doi.org/10.1641/0006-3568\(2000\)050%255B0188:TPONTP%255D2.3.CO;2](https://doi.org/10.1641/0006-3568(2000)050%255B0188:TPONTP%255D2.3.CO;2)
- Vander Mijnsbrugge, K., Bischoff, A. et Smith, B. (2010). A question of origin: Where and how to collect seed for ecological restoration. *Basic and Applied Ecology*, 11(4), 300-311. <https://doi.org/10.1016/j.baae.2009.09.002>
- Vivian-Smith, G. et Handel, S. N. (1996). Freshwater wetland restoration of an abandoned sand mine: Seed bank recruitment dynamics and plant colonization. *Wetlands*, 16(2), 185-196. <https://doi.org/10.1007/BF03160692>
- White, A., Fant, J. B., Havens, K., Skinner, M. & Kramer, A. T. (2018). Restoring species diversity: Assessing capacity in the U.S. native plant industry. *Restoration Ecology*, 26(4), 605–611. <https://doi.org/10.1111/rec.12705>
- Wilson, S. D., Moore, D. R. J. et Keddy, P. A. (1993). Relationships of marsh seed banks to vegetation patterns along environmental gradients. *Freshwater Biology*, 29(3), 361-370. <https://doi.org/10.1111/j.1365-2427.1993.tb00770.x>
- Yakimowski, S. B., Hager, H. A. et Eckert, C. G. (2005). Limits and effects of invasion by the nonindigenous wetland plant *Lythrum salicaria* (purple loosestrife): a seed bank analysis. *Biological Invasions*, 7(4), 687-698. <https://doi.org/10.1007/s10530-004-5858-y>

Yetka, L. A. et Galatowitsch, S. M. (1999). Factors Affecting Revegetation of *Carex lacustris* and *Carex stricta* from Rhizomes. *Restoration Ecology*, 7(2), 162-171.  
<https://doi.org/10.1046/j.1526-100X.1999.72008.x>

Zinnen, J., Broadhurst, L. M., Gibson-Roy, P., Jones, T. A. et Matthews, J. W. (2021). Seed production areas are crucial to conservation outcomes: benefits and risks of an emerging restoration tool. *Biodiversity and Conservation*, 30(5), 1233-1256.  
<https://doi.org/10.1007/s10531-021-02149-z>